

Hoffman's Limit Theorem

Akihiro Munemasa

Graduate School of Information Sciences
Tohoku University

joint work with A. Gavriluk, Y. Sano, T. Taniguchi
August 20, 2019

The International Conference and PhD-Master Summer School on
Groups and Graphs, Designs and Dynamics
Yichang, China

History (λ_{\min} = smallest eigenvalue)

$$\lim_{t \rightarrow \infty} \lambda_{\min} \begin{bmatrix} A & C \otimes \mathbf{1}_t \\ C^\top \otimes \mathbf{1}_t^\top & I \otimes (J_t - I_t) \end{bmatrix} = \lambda_{\min}(A - CC^\top).$$

- Hoffman (SIAM, 1969) stated a theorem (Hoffman's limit theorem), "is shown in [4]" where [4]=Hoffman & Ostrowski, "to appear" was never published.
- Hoffman (LAA, 1977), citing above, proved a **theorem about graphs with $\lambda_{\min} \in (-2, -1)$** and $\lambda_{\min} \in (-1 - \sqrt{2}, -2)$.
- Jang–Koolen–M.–Taniguchi (AMC, 2014) gave a graph theoretic proof.
- Hoffman (Geom. Ded. 1977), proved **signed** graph version of the limit theorem.

Today, we give a **Hermitian matrix** version of the limit theorem and an application to **signed graphs with $\lambda_{\min} \in (-2, -1)$** .

What is the spectrum of a graph

The **spectrum** of a graph means the multiset of eigenvalues of its adjacency matrix.

$$\text{Spec} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = \{1, -1\},$$

$$\text{Spec}(K_n) = \text{Spec}(J_n - I_n) = \{[n-1]^1, [-1]^{n-1}\},$$

$$\text{Spec} \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} = \{\sqrt{2}, 0, -\sqrt{2}\},$$

$$\text{Spec} \begin{bmatrix} 0 & 0 & \mathbf{1}_t \\ 0 & 0 & \mathbf{1}_t \\ \mathbf{1}_t^\top & \mathbf{1}_t^\top & J_t - I_t \end{bmatrix} = ?$$

→ on blackboard

The smallest eigenvalue of a graph

Denote by $\lambda_{\min}(\cdot)$ the smallest eigenvalue of a matrix or a graph.

$$\lambda_{\min} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = -1,$$

$$\lambda_{\min}(K_n) = \lambda_{\min}(J_n - I_n) = -1,$$

$$\lambda_{\min} \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} = -\sqrt{2},$$

$$\lambda_{\min} \begin{bmatrix} 0 & 0 & \mathbf{1}_t \\ 0 & 0 & \mathbf{1}_t \\ \mathbf{1}_t^\top & \mathbf{1}_t^\top & J_t - I_t \end{bmatrix} = ?$$

Γ is connected and $\lambda_{\min}(\Gamma) = -1 \implies \Gamma \cong K_n$.

$$\text{Spec} \begin{bmatrix} 0 & 0 & \mathbf{1}_t \\ 0 & 0 & \mathbf{1}_t \\ \mathbf{1}_t^\top & \mathbf{1}_t^\top & J_t - I_t \end{bmatrix} = \text{Spec} \begin{bmatrix} 0 & 0 & t \\ 0 & 0 & t \\ 1 & 1 & t-1 \end{bmatrix} \cup \text{Spec}(-I_t).$$

$$\begin{aligned} \lambda_{\min} \begin{bmatrix} 0 & 0 & \mathbf{1}_t \\ 0 & 0 & \mathbf{1}_t \\ \mathbf{1}_t^\top & \mathbf{1}_t^\top & J_t - I_t \end{bmatrix} &= \lambda_{\min} \begin{bmatrix} 0 & 0 & t \\ 0 & 0 & t \\ 1 & 1 & t-1 \end{bmatrix} \\ &= \min\{z \mid z(z^2 - (t-1)z - 2t) = 0\} \\ &= \frac{t-1 - \sqrt{t^2 + 6t + 1}}{2} \rightarrow -2 \quad (t \rightarrow \infty). \end{aligned}$$

Shortcut (?)

$$\min\{z \mid (z+2) - \frac{1}{t}(z^2+z) = 0\} \rightarrow \min\{z \mid z+2 = 0\} = -2.$$

Hurwitz's theorem

Rahman & Schmeisser, "Analytic Theory of Polynomials,"
Theorem 1.3.8

Theorem

Let $(f_t)_{t=1}^{\infty}$ be a sequence of analytic functions defined in a region $\Omega \subseteq \mathbb{C}$. Suppose

$$f_t \rightarrow f \neq 0 \quad (t \rightarrow \infty)$$

uniformly on every compact subset of Ω . Then for $\zeta \in \Omega$, the following are equivalent:

- 1 ζ is a zero of f of multiplicity m ,
- 2 $\zeta \in \exists U \subseteq \Omega$ (neighbourhood), $\forall \varepsilon > 0$, $\exists n_0 < \forall t$, f_t has exactly m zeros in the ε -neighbourhood of ζ .

Theorem (Hoffman's limit theorem)

Let

$$\begin{bmatrix} A & C \\ C^\top & 0 \end{bmatrix}$$

be the adjacency matrix of a graph. Then

$$\lim_{t \rightarrow \infty} \lambda_{\min} \begin{bmatrix} A & C \otimes \mathbf{1}_t \\ C^\top \otimes \mathbf{1}_t^\top & I \otimes (J_t - I_t) \end{bmatrix} = \lambda_{\min}(A - CC^\top).$$

→ on blackboard

$$A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, C = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, A - CC^\top = - \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix},$$

which has $\lambda_{\min} = -2$. Note

$$\lambda_{\min} \begin{bmatrix} 0 & 0 & \mathbf{1}_t \\ 0 & 0 & \mathbf{1}_t \\ \mathbf{1}_t^\top & \mathbf{1}_t^\top & J_t - I_t \end{bmatrix} > -2.$$

Proof of Hoffman's limit theorem

$$\lim_{t \rightarrow \infty} \lambda_{\min} \begin{bmatrix} A & C \otimes \mathbf{1}_t \\ C^T \otimes \mathbf{1}_t^T & I \otimes (J_t - I_t) \end{bmatrix} = \lim_{t \rightarrow \infty} \lambda_{\min} \begin{bmatrix} A & tC \\ C^T & (t-1)I \end{bmatrix}$$

Since

→ on blackboard

$$\left| zI - \begin{bmatrix} A & tC \\ C^T & (t-1)I \end{bmatrix} \right| = t \left| A - CC^T - zI + \frac{z+1}{t}(zI - A) \right|,$$

the spectrum containing $\lambda_{\min} \rightarrow \text{Spec}(A - CC^T)$.

$\lambda_{\min} \rightarrow \lambda_{\min}(A - CC^T)$, proving the theorem.

The same proof shows the Hermitian matrix version:

Theorem

Let

$$\begin{bmatrix} A & C \\ C^* & 0 \end{bmatrix}$$

be a Hermitian matrix, and let D be a positive definite Hermitian matrix. Then

$$\lim_{t \rightarrow \infty} \lambda_{\min} \begin{bmatrix} A & C \otimes \mathbf{1}_t \\ C^* \otimes \mathbf{1}_t^\top & D \otimes (J_t - I_t) \end{bmatrix} = \lambda_{\min}(A - CD^{-1}C^*).$$

A **signed graph** is a graph with edge weight $+1$ or -1 . The adjacency matrix is then a $(0, \pm 1)$ matrix.

- Switching equivalence = conjugation by a $(0, \pm 1)$ monomial matrix
- $\delta(G) :=$ minimum degree of G .

Theorem

There exists a function $f : (-2, -1) \rightarrow \mathbb{R}$ such that, for each $\lambda \in (-2, -1)$, if G is a connected **signed** graph with $\lambda_{\min}(G) \geq \lambda$, $\delta(G) \geq f(\lambda)$, then G is switching equivalent to a complete graph.

The proof is a simplification of Hoffman's original by incorporating Cameron–Goethals–Seidel–Shult (1976), Greaves–Koolen–M.–Sano–Taniguchi (2015).

Proof (part 1)

Fix $\lambda \in (-2, -1)$. To prove this theorem, it suffices to show that,

$$\begin{array}{l} \lambda_{\min}(G) \geq \lambda \\ \delta(G) \text{ sufficiently large} \end{array} \implies G \text{ is sw. eq. } K_n.$$

By Cameron–Goethals–Seidel–Shult (1976), we may assume G is represented by A_m or D_m (ignoring E_6, E_7, E_8).

But $A_m \subseteq D_{m+1}$, so

Proof (part 2)

$$\begin{aligned} \lambda_{\min}(G) &\geq \lambda \\ \delta(G) \text{ sufficiently large} &\implies G \text{ is sw. eq. } K_n. \\ G \text{ is represented by } D_m & \end{aligned}$$

Greaves–Koolen–M.–Sano–Taniguchi (2015) classified such signed graphs. In particular,

Theorem

Let G be a connected **signed** graph represented by D_m and $\lambda_{\min}(G) > -2$. Then there exists a tree T such that the line graph $L(T)$ of T is switching equivalent to G with possibly one vertex removed.

Here we illustrate the proof when $G - u$ is sw. eq. to $L(T)$.

→ on blackboard

Recall the **Hermitian** adjacency matrix $H = H(\Delta)$ of a digraph Δ :

$$H_{xy} = \begin{cases} 1 & \text{if } x \rightleftarrows y \\ i & \text{if } x \rightarrow y \\ -i & \text{if } x \leftarrow y \\ 0 & \text{otherwise} \end{cases}$$

Theorem

There exists a function $f : (-2, -1) \rightarrow \mathbb{R}$ such that, for each $\lambda \in (-2, -1)$, if Δ is a connected digraph with $\lambda_{\min}(H(\Delta)) \geq \lambda$, $\delta(\overline{\Delta}) \geq f(\lambda)$, then Δ is **switching equivalent** to a complete graph.

- Switching equivalence = conjugation by a $(0, \pm 1, \pm i)$ monomial matrix, and possibly taking the transpose
- $\delta(\overline{\Delta}) :=$ minimum degree of the underlying undirected graph of Δ .

Theorem

There exists a function $f : (-2, -1) \rightarrow \mathbb{R}$ such that, for each $\lambda \in (-2, -1)$,

- 1 for connected **signed** graph G , $\lambda_{\min}(G) \geq \lambda$,
 $\delta(G) \geq f(\lambda) \implies G$ sw. eq. K_n .
- 2 for connected **digraph** Δ , $\lambda_{\min}(H(\Delta)) \geq \lambda$,
 $\delta(\overline{\Delta}) \geq f(\lambda) \implies \Delta$ sw. eq. K_n .

The digraph version is immediate from signed graph version by considering the **associated** signed graph:

$$H(\Delta) = A + iB \quad (A = A^\top, B = -B^\top) \implies A(G) = \begin{bmatrix} A & B \\ B^\top & A \end{bmatrix}$$

- $\text{Spec } H(\Delta)^{\times 2} = \text{Spec } G$, so $\lambda_{\min} H(\Delta) = \lambda_{\min} G$.
- $\delta(\overline{\Delta}) = \delta(G)$.

Further results yet to be generalized: Hoffman (1977):
 $(-1 - \sqrt{2}, -2)$, Woo & Neumaier (1995).

The idea of associated signed graph comes from

- **Masaaki Kitazume** and A. M., Even unimodular Gaussian lattices of rank 12, J. Number Theory (2002).

Gaussian lattices of rank 12 \leftrightarrow Euclidean lattices of rank 24

A digraph with n vertices \rightarrow its associated signed graph has $2n$ vertices:

$$H(\Delta) = A + iB \quad (A = A^\top, B = -B^\top) \implies A(G) = \begin{bmatrix} A & B \\ B^\top & A \end{bmatrix}$$

Given a signed adjacency matrix S of order $2n$, find a hermitian matrix $H = A + iB$ of order n such that S is switching equivalent to

$$\begin{bmatrix} A & B \\ B^\top & A \end{bmatrix}$$